

Other Publications:

- Tan, H. M., Wang, X., Soh, S. F., Tan, S., Zhao, J., Yong, E. L., Lee, H. K., and Gong, Y. (2012). Preparation and Application of Mixed Octadecylsilyl- and (3-(C-Methylcalix[4]Resorcinarene)-Hydroxypropoxy)-Propylsilyl-Appended Silica Particles as Stationary Phase for High-Performance Liquid Chromatography. *Instrumentation Science & Technology*, 40(2-3), 100-111. doi: 10.1080/10739149.2011.651674
- Tan, H. M., Soh, S. F., Zhao, J., Yong, E. L., and Gong, Y. (2011). Preparation and application of methylcalix[4]resorcinarene-bonded silica particles as chiral stationary phase in high-performance liquid chromatography. *Chirality*, 23 Suppl 1, E91-97. doi: 10.1002/chir.20983
- Kang, J., Shin, J., Cho, D., Jeong, Y. Park, C., Soh, S.F., Lai, C.S.,and Tiekink, E.R.T (2010). Steric Control over Supramolecular Polymer Formation in trans-1,2-Bis(4-pyridyl)ethylene Adducts of Zinc Xanthates: Implications for Luminescence Cryst. Growth Des., 10 (3), pp 1247–1256 DOI: 10.1021/cg9012735
- Soh, S.F., Lai, C.S. and Tiekink, E.R.T (2002). (4,7-Dimethyl-1,10-phenanthroline)-bis(O-isopropylidithiocarbonato)zinc(II) *Acta Crystallographica Section E Structure Reports Online*. 11/2002; 58(11). DOI: 10.1107/S1600536802018433